arXiv:1204.2133v1 [math.NT] 10 Apr 2012

TRIVIAL INTEGRAL GALOIS MODULE STRUCTURE OF WEAKLY RAMIFIED EXTENSIONS OF P-ADIC FIELDS
PRELIMINARY VERSION

HENRI JOHNSTON

Abstract. Let p be a rational prime and let K be a finite extension of $\mathbb{Q}_p$. Let L be a finite Galois extension extension of K and let $G = \mathrm{Gal}(L/K)$. We say that L/K is weakly ramified if and only if its second ramification group is trivial. Let $\mathcal{O}_L$ be the valuation ring of L and let $\mathfrak{P}_L$ be its maximal ideal. We show that if L/K is weakly ramified then $\mathfrak{P}_L$ is free over the group ring $\mathcal{O}_K[G]$, and from this we deduce that if L/K is wildly and weakly ramified then $\mathcal{O}_L$ is free over its associated order in the group algebra $K[G]$.

1. Introduction

Let p be a rational prime and let K be a finite extension of $\mathbb{Q}_p$ with valuation ring $\mathcal{O}_K$. Let L be a finite Galois extension extension of K and let $G = \mathrm{Gal}(L/K)$. Let $\mathcal{O}_L$ be the valuation ring of L and let $\mathfrak{P}_L$ be its maximal ideal. We shall be concerned with the structure of both $\mathfrak{P}_L$ over the group ring $\mathcal{O}_K[G]$ and of $\mathcal{O}_L$ over its associated order

$$\mathcal{A}_{L/K} = \{x \in K[G] \mid x\mathcal{O}_L \subseteq \mathcal{O}_L\}.$$

We recall that the ramification groups of L/K (in the lower numbering) are the groups

$$G_i = \{\sigma \in G \mid (\sigma - 1)(\mathcal{O}_L) \subseteq \mathfrak{P}_L^{i+1}\} \quad \text{for} \quad i \geq -1.$$

Thus L/K is unramified if and only if G_0 is trivial and is tamely ramified if and only if G_1 is trivial. We say that L/K is weakly ramified if and only if G_2 is trivial.

It is well-known that $\mathcal{A}_{L/K}$ coincides with the group ring $\mathcal{O}_K[G]$ precisely when L/K is tamely ramified, and that in this case $\mathcal{O}_L$ is free as a module over $\mathcal{O}_K[G]$ (for a proof of the second assertion see [Kaw86]). Furthermore, if L/K is tamely ramified then [Ull70, Theorem 1] shows that every fractional ideal of $\mathcal{O}_L$ is free as a module over $\mathcal{O}_K[G]$. In the case that L/K is totally and weakly ramified, [Ull70, Theorem 2] shows that $\mathfrak{P}_L$ is free over $\mathcal{O}_K[G]$; from this it is straightforward to deduce that $\mathcal{O}_L$ is free over $\mathcal{A}_{L/K}$ and that $\mathcal{A}_{L/K} = \mathcal{O}_K[G][\pi_K^{-1}\mathrm{Tr}_G]$ where π_K is any prime element of K and $\mathrm{Tr}_G = \sum_{\tau \in G} \tau$. If L/K is abelian and weakly ramified (but not necessarily totally ramified), then [Byo99, Theorem 5] shows that $\mathcal{O}_L$ is free over $\mathcal{A}_{L/K}$ and that $\mathcal{A}_{L/K} = \mathcal{O}_K[G][\pi_K^{-1}\mathrm{Tr}_{G_0}]$. In the present article, we generalise these results as follows.

Theorem 1.1. *Let p be a rational prime and let K be a finite extension of $\mathbb{Q}_p$. Let L/K be a finite Galois extension that is weakly ramified and let $G = \mathrm{Gal}(L/K)$. Then there exists $\varepsilon \in \mathfrak{P}_L \backslash \mathfrak{P}_L^2$ such that ε is a free generator of $\mathfrak{P}_L$ over $\mathcal{O}_K[G]$.*

Date: Version of 5th April 2012.

2010 *Mathematics Subject Classification.* 11R33.

Key words and phrases. Weakly ramified, local fields,

Corollary 1.2. *Assume the situation and notation of Theorem 1.1. Let* π_K *be any prime element of* K *and let* $\mathrm{Tr}_{G_0} = \sum_{\tau \in G_0} \tau$. *If* L/K *is wildly and weakly ramified then* $\mathcal{A}_{L/K} = \mathcal{O}_K[G][\pi_K^{-1}\mathrm{Tr}_{G_0}]$ *and* ε *is a free generator of* $\mathcal{O}_L$ *over* $\mathcal{A}_{L/K}$.

2. INGREDIENTS

In this section we review several results that form the 'ingredients' of the proof of the main results of the present article. By a 'local field' we mean a complete field under a discrete valuation with a finite residue field.

2.1. **Totally and weakly ramified p-extensions.**

Proposition 2.1. *Let* p *be a rational prime and let* K *be a finite extension of* $\mathbb{Q}_p$. *Let* L/K *be a totally and weakly ramified finite* p*-extension and let* $G = \mathrm{Gal}(L/K)$. *Then any prime element* π_L *of* L *is a free generator of* $\mathfrak{P}_L$ *over* $\mathcal{O}_K[G]$.

Remark 2.2. The hypotheses force G to be an elementary abelian p-group.

Proof. This result is proven in [Vos81, Proposition 2]. Alternatively: by [Byo99, Corollary 4.3] any prime element π_L in L is a free generator of $\mathcal{O}_L$ over $\mathcal{O}_K[G][\pi_K^{-1}\mathrm{Tr}_G]$ where $\mathrm{Tr}_G = \sum_{\tau \in G} \tau$, and from this [Vin05, Proposition 2.4] shows that any π_L is a free generator of $\mathfrak{P}_L$ over $\mathcal{O}_K[G]$. Note that under the more general hypothesis that L/K is totally and weakly ramified (but not necessarily a p-extension), [Ull70, Theorem 2] shows that $\mathfrak{P}_L$ is free over $\mathcal{O}_K[G]$, though a generator is not specified. ☐

2.2. **Totally and tamely ramified extensions.**

Lemma 2.3. *Let* L/K *be a totally and tamely ramified extension of local fields of degree* e. *Then there exist prime elements* π_L *and* π_K *in* L *and* K *respectively such that* $\pi_L^e = \pi_K$.

Proof. See [Lan94, Chapter 2, Proposition 12], for example. ☐

Proposition 2.4. *Let* L/K *be a totally and tamely ramified Galois extension of local fields of degree* e *and let* $G = \mathrm{Gal}(L/K)$. *Let* π_L *be as in Lemma 2.3 and let*

$$\alpha = 1 + \pi_L + \cdots + \pi_L^{e-1}.$$

If $I = \mathfrak{P}_L^{v(I)}$ *is any fractional ideal of* $\mathcal{O}_L$ *then* $\pi_L^{v(I)}\alpha$ *is a free generator of* I *over* $\mathcal{O}_K[G]$.

Proof. As noted in [Ere91, §7.1]: the straightforward determinant calculation given in [Kaw86] shows that α is a free generator of $\mathcal{O}_L$ over $\mathcal{O}_K[G]$, and from this the desired result follows. ☐

Remark 2.5. For any choice of $u_0, \ldots, u_{e-1} \in \mathcal{O}_K^\times$, we can in fact take $\alpha = \sum_{i=0}^{e-1} u_i \pi_L^i$.

2.3. **Unramified extensions.**

Proposition 2.6. *Let* L/K *be a finite unramified extension of local fields and let* $G = \mathrm{Gal}(L/K)$. *Then there exists* $\beta \in \mathcal{O}_L$ *such that* β *is a free generator of* $\mathcal{O}_L$ *over* $\mathcal{O}_K[G]$.

Proof. This is standard (see [Kaw86], for example), but we give a proof as it is short. Let $\overline{L}$ and $\overline{K}$ be the residue fields of L and K, respectively. As L/K is unramified, G identifies with $\mathrm{Gal}(\overline{L}/\overline{K})$. By the Normal Basis Theorem, there exists $\overline{\beta} \in \overline{L}$ such that $\overline{L} = \overline{K}[G] \cdot \overline{\beta}$. Let β be any lift of $\overline{\beta}$ to $\mathcal{O}_L$. Then a straightforward application of Nakayama's Lemma gives $\mathcal{O}_L = \mathcal{O}_K[G] \cdot \beta$. Thus the map $\mathcal{O}_K[G] \longrightarrow \mathcal{O}_L$ given by $x \mapsto x \cdot \beta$ is surjective and a rank argument shows that it is also injective. ☐

2.4. **A splitting lemma.**

Lemma 2.7. *Let L/K be a finite Galois extension of local fields. Let K'/K be the unique unramified extension of degree $[L:K]$. Let $L' = LK'$. Then L'/K is Galois. Let $G = \mathrm{Gal}(L'/K)$. Then $G_0 = \mathrm{Gal}(L'/K')$ is the inertia subgroup of G, and G decomposes as a semi-direct product $G = G_0 \rtimes H_0$ for some cyclic subgroup H_0 of G.*

Proof. The proof given here is an adaptation of that of [Let98, Lemma 1]. Since L/K and K'/K are both Galois, so is L'/K. By considering ramification degrees, it is straightforward to check that $\mathrm{Gal}(L'/K')$ is the inertia subgroup of $\mathrm{Gal}(L'/K)$. Consider the exact sequence

$$1 \longrightarrow \mathrm{Gal}(L'/K') \longrightarrow \mathrm{Gal}(L'/K) \xrightarrow{\rho} \mathrm{Gal}(K'/K) \longrightarrow 1.$$

Let $\sigma \in \mathrm{Gal}(K'/K)$ be the Frobenius element (or indeed any generator of this cyclic group) and take any $\tau \in \mathrm{Gal}(L'/K)$ with $\rho(\tau) = \sigma$. Let $d = [L:K] = [K':K]$. Then τ^d is the identity on both L and K', so we have $\tau^d = \mathrm{id}_{L'}$. Therefore $\varphi : \mathrm{Gal}(K'/K) \longrightarrow \mathrm{Gal}(L'/K)$, defined by $\varphi(\sigma) = \tau$, is a splitting homomorphism for the above exact sequence. Hence we may take $H_0 = \varphi(\mathrm{Gal}(K'/K))$. $\square$

3. TOTALLY AND WEAKLY RAMIFIED EXTENSIONS

Let p be a rational prime and let M be a finite extension of $\mathbb{Q}_p$. Let L/M be a totally and weakly ramified finite Galois extension and let $G_0 = \mathrm{Gal}(L/M)$. Since G_2 is trivial, G_1 is an elementary abelian p-group. Furthermore, G_0 decomposes as a semi-direct product $G_0 = G_1 \rtimes H_1$ for some cyclic subgroup H_1 of G_0. Let $E = L^{G_1}$ and $F = L^{H_1}$ be the subfields of L fixed by G_1 and H_1, respectively. Note that the choice of H_1 (and hence of F) is not necessarily unique and that the order of H_1 is prime to p. We identify $\mathrm{Gal}(E/M)$ with $H_1 = \mathrm{Gal}(L/F)$ via the restriction map $H_1 \to \mathrm{Gal}(E/K)$, $\gamma \mapsto \gamma|_E$. The situation is represented by the following field diagram.



Both L/E and F/M are are totally and wildly ramified p-extensions and both L/F and E/M are totally and tamely ramified. Note that F/M need not be Galois.

Define r by $p^r = [L:E] = [F:M]$ and let $s = [L:F] = [E:M]$. Since E/M is totally and tamely ramified, by Lemma 2.3 there exist prime elements π_E and π_M of E and M respectively such that $\pi_E^s = \pi_M$. By Bezout's Theorem, there exist integers a, b such that $ap^r + bs = 1$. Let π_F be any prime element of F.

Proposition 3.1. *We have $\mathfrak{P}_L = \mathcal{O}_M[G_0] \cdot (\pi_F^b \pi_E^a \alpha)$ where $\alpha = 1 + \pi_E + \cdots + \pi_E^{s-1}$.*

Proof. Write $G_1 = \{\tau_i\}$ and $H_1 = \{\sigma_j\}$. Since L/M is weakly ramified, it follows directly from the definition of the ramification groups that L/E is also weakly ramified. Hence by Proposition 2.1 any any prime element π_L in L is a basis of $\mathfrak{P}_L$ over $\mathcal{O}_E[G_1]$. However, if v_L is the normalised valuation on L then we have $v_L(\pi_F^b \pi_E^a) = bs + ap^r = 1$, and so in particular we may take $\pi_L = \pi_F^b \pi_E^a$. Furthermore, by Proposition 2.4 we have that $\pi_E^a \alpha$

is a free generator of the fractional ideal $\pi_E^a\mathcal{O}_E$ over $\mathcal{O}_M[H_1]$. Therefore we have

$$\begin{aligned}
\mathfrak{P}_L = \mathcal{O}_E[G_1]\cdot(\pi_F^b\pi_E^a) &= \bigoplus_i \tau_i(\pi_F^b\pi_E^a)\mathcal{O}_E \\
&= \bigoplus_i \tau_i(\pi_F^b)(\pi_E^a\mathcal{O}_E) \quad \text{since } \pi_E \in E = L^{G_1} \\
&= \bigoplus_i \tau_i(\pi_F^b)(\mathcal{O}_M[H_1]\cdot\pi_E^a\alpha) = \bigoplus_i \tau_i(\pi_F^b)\bigoplus_j \sigma_j(\pi_E^a\alpha)\mathcal{O}_M \\
&= \bigoplus_i\bigoplus_j \tau_i(\pi_F^b)\sigma_j(\pi_E^a\alpha)\mathcal{O}_M \\
&= \bigoplus_i\bigoplus_j \tau_i\sigma_j(\pi_F^b)\sigma_j(\pi_E^a\alpha)\mathcal{O}_M \quad \text{since } \pi_F \in F = L^{H_1} \\
&= \bigoplus_i\bigoplus_j \tau_i\sigma_j(\pi_F^b)\tau_i\sigma_j(\pi_E^a\alpha)\mathcal{O}_M \quad \text{since } \sigma_j(\pi_E^a\alpha) \in E = L^{G_1} \\
&= \bigoplus_i\bigoplus_j \tau_i\sigma_j(\pi_F^b\pi_E^a\alpha)\mathcal{O}_M = \mathcal{O}_M[G_0]\cdot(\pi_F^b\pi_E^a\alpha). \qquad \square
\end{aligned}$$

4. Split weakly ramified extensions

Let p be a rational prime and let K be a finite extension of $\mathbb{Q}_p$. Let L/K be a weakly ramified finite Galois extension and let $G = \mathrm{Gal}(L/K)$. Suppose that $G = G_{-1}$ decomposes as a semi-direct product $G = G_0 \rtimes H_0$ for some subgroup H_0 of G, where H_0 is necessarily cyclic. Let $M = L^{G_0}$ be the inertia subfield and let $N = L^{H_0}$. Note that the choice of H_0 (and hence of N) is not necessarily unique. We identify $\mathrm{Gal}(M/K)$ with $H_0 = \mathrm{Gal}(L/N)$ via the restriction map $H_0 \to \mathrm{Gal}(M/K)$, $\gamma \mapsto \gamma|_M$. The extension L/M 'decomposes' exactly as in §3 and we henceforth assume all the notation used therein. The situation is represented by the following pair of field diagrams.





We combine these into the following field diagram.



Here we have identified H_0 with the Galois group of the relevant extension as above, and unmarked extensions are not necessarily Galois.

We now choose elements in the various intermediate fields, from which we will construct a free generator of $\mathfrak{P}_L$ over $\mathcal{O}_K[G]$. Since $F/F\cap N$ is unramified, we may choose π_F to be a prime element of $F \cap N$. As $E \cap N/K$ is totally and tamely ramified, by Lemma 2.3 there exist prime elements $\pi_{E\cap N}$ and π_K such that $\pi_{E\cap N}^s = \pi_K$. Since both $E/E \cap N$ and M/K are unramified, we may take $\pi_E = \pi_{E\cap N}$ and $\pi_M = \pi_K$. Therefore all the chosen prime elements belong to N. Finally, since M/K is unramified, by Proposition 2.6 there exists $\beta \in \mathcal{O}_M$ such that $\mathcal{O}_M = \mathcal{O}_K[H_0] \cdot \beta$.

Proposition 4.1. *We have* $\mathfrak{P}_L = \mathcal{O}_K[G] \cdot (\pi_F^b \pi_E^a \alpha\beta)$ *where* $\alpha = 1 + \pi_E + \cdots + \pi_E^{s-1}$.

Proof. Let $\gamma = \pi_F^b \pi_E^a \alpha$. Then by Proposition 3.1 we have $\mathfrak{P}_L = \mathcal{O}_M[G_0] \cdot \gamma$. A key point is that γ belongs to N since all the primes elements above were chosen to be in N. Write $G_0 = \{\tau_i\}$ and $H_0 = \{\sigma_j\}$. Then we have

$$\begin{aligned}
\mathfrak{P}_L = \mathcal{O}_M[G_0] \cdot \gamma = \bigoplus_i \tau_i(\gamma)\mathcal{O}_M &= \bigoplus_i \tau_i(\gamma)(\mathcal{O}_K[H_0] \cdot \beta) \\
&= \bigoplus_i \tau_i(\gamma) \bigoplus_j \sigma_j(\beta)\mathcal{O}_K = \bigoplus_i \bigoplus_j \tau_i(\gamma)\sigma_j(\beta)\mathcal{O}_K \\
&= \bigoplus_i \bigoplus_j \tau_i\sigma_j(\gamma)\sigma_j(\beta)\mathcal{O}_K \quad \text{since } \gamma \in N = L^{H_0} \\
&= \bigoplus_i \bigoplus_j \tau_i\sigma_j(\gamma)\tau_i\sigma_j(\beta)\mathcal{O}_K \quad \text{since } \sigma_j(\beta) \in M = L^{G_0} \\
&= \bigoplus_i \bigoplus_j \tau_i\sigma_j(\gamma\beta)\mathcal{O}_K = \mathcal{O}_K[G] \cdot (\gamma\beta). \qquad \square
\end{aligned}$$

5. Proofs of Theorem 1.1 and Corollary 1.2

Proof of Theorem 1.1. Let K'/K be the unique unramified extension of degree $[L : K]$, and set $L' = LK'$. Since L'/L is unramified, [Byo99, Proposition 4.4] implies that L'/K is weakly ramified. Now Lemma 2.7 shows that L'/K is of the form considered in §4. Hence by Proposition 4.1 there exists an $\varepsilon' \in \mathcal{O}_{L'}$ such that $\mathfrak{P}_{L'} = \mathcal{O}_K[\mathrm{Gal}(L'/K)] \cdot \varepsilon'$. As L'/L is unramified, [FT93, Corollary to Theorem 26] shows that $\mathrm{Tr}_{L'/L}(\mathfrak{P}_{L'}) = \mathfrak{P}_L$. Thus we obtain

$$\mathfrak{P}_L = \mathrm{Tr}_{L'/L}(\mathcal{O}_K[\mathrm{Gal}(L'/K)] \cdot \varepsilon') = \mathcal{O}_K[\mathrm{Gal}(L'/K)] \cdot \mathrm{Tr}_{L'/L}(\varepsilon') = \mathcal{O}_K[G] \cdot \mathrm{Tr}_{L'/L}(\varepsilon').$$

Hence the map $\mathcal{O}_K[G] \longrightarrow \mathfrak{P}_L$ given by $x \mapsto x \cdot \mathrm{Tr}_{L'/L}(\varepsilon')$ is surjective and a rank argument shows that it is also injective. Therefore we may take $\varepsilon = \mathrm{Tr}_{L'/L}(\varepsilon')$. It is straightforward to see that we must have $\varepsilon \in \mathfrak{P}_L \backslash \mathfrak{P}_L^2$. $\square$

Proof of Corollary 1.2. Let $F = L^{G_0}$ be the inertia subfield of L. Since L/F is wildly ramified, we have $\mathrm{Tr}_{L/F}(\mathcal{O}_L) \subseteq \mathfrak{P}_F$ by [FT93, Theorem 26 (b)]. Since F/K is unramified, we hence have $\pi_K^{-1}\mathrm{Tr}_{L/F}(\mathcal{O}_L) \subseteq \pi_K^{-1}\mathfrak{P}_F = \mathcal{O}_F$. Therefore $\mathcal{O}_K[G][\pi_K^{-1}\mathrm{Tr}_{G_0}] \subseteq \mathcal{A}_{L/K}$.

Suppose that there exists $\varepsilon \in \mathcal{O}_L$ such that $\mathfrak{P}_L = \mathcal{O}_K[G] \cdot \varepsilon$. Then we have

$$\mathcal{O}_K[G][\pi_K^{-1}\mathrm{Tr}_{G_0}] \cdot \varepsilon \subseteq \mathcal{A}_{L/K} \cdot \varepsilon \subseteq \mathcal{O}_L.$$

Considering generalised module indices (see [FT93, II.4] for background), we have

$$\begin{aligned}
[\mathcal{O}_L : \mathfrak{P}_L]_{\mathcal{O}_K} = [\mathcal{O}_F : \mathfrak{P}_F]_{\mathcal{O}_K} = \mathfrak{P}_K^{[F:K]} &= [\mathcal{O}_K[G][\pi_K^{-1}\mathrm{Tr}_{G_0}] : \mathcal{O}_K[G]]_{\mathcal{O}_K} \\
&= [\mathcal{O}_K[G][\pi_K^{-1}\mathrm{Tr}_{G_0}] \cdot \varepsilon : \mathcal{O}_K[G] \cdot \varepsilon]_{\mathcal{O}_K} = [\mathcal{O}_K[G][\pi_K^{-1}\mathrm{Tr}_{G_0}] \cdot \varepsilon : \mathfrak{P}_L]_{\mathcal{O}_K}.
\end{aligned}$$

Therefore we have $\mathcal{O}_K[G][\pi_K^{-1}\mathrm{Tr}_{G_0}] \cdot \varepsilon = \mathcal{O}_L$. Hence the map $\mathcal{O}_K[G][\pi_K^{-1}\mathrm{Tr}_{G_0}] \longrightarrow \mathcal{O}_L$ given by $x \mapsto x \cdot \varepsilon$ is surjective and a rank argument shows that it is also injective. Thus

ε is a free generator of $\mathcal{O}_L$ over $\mathcal{O}_K[G][\pi_K^{-1}\mathrm{Tr}_{G_0}]$ and so it follows that

$$\mathcal{A}_{L/K} = \mathcal{O}_K[G][\pi_K^{-1}\mathrm{Tr}_{G_0}]. \qquad \square$$

Remark 5.1. Let L/K be any finite tamely ramified Galois extension of local fields and let $G = \mathrm{Gal}(L/K)$. The results of Propositions 2.4 and 2.6 and Lemma 2.7 can be combined in a similar way to the proofs of Theorem 1.1 and Proposition 4.1 to recover the well-known result that there exists $\varepsilon \in \mathcal{O}_L$ such that ε is a free generator of $\mathcal{O}_L$ over $\mathcal{O}_K[G]$ in this case. This is essentially the approach taken in [Kaw86].

REFERENCES

[Byo99] N. P. Byott, *Integral Galois module structure of some Lubin-Tate extensions*, J. Number Theory **77** (1999), no. 2, 252–273. MR 1702149 (2000f:11156)

[Ere91] B. Erez, *The Galois structure of the square root of the inverse different*, Math. Z. **208** (1991), no. 2, 239–255. MR 1128708 (92g:11108)

[FT93] A. Fröhlich and M. J. Taylor, *Algebraic number theory*, Cambridge Studies in Advanced Mathematics, vol. 27, Cambridge University Press, Cambridge, 1993. MR 1215934 (94d:11078)

[Kaw86] F. Kawamoto, *On normal integral bases of local fields*, J. Algebra **98** (1986), no. 1, 197–199. MR 825142 (87e:11137)

[Lan94] S. Lang, *Algebraic number theory*, second ed., Graduate Texts in Mathematics, vol. 110, Springer-Verlag, New York, 1994. MR 1282723 (95f:11085)

[Let98] G. Lettl, *Relative Galois module structure of integers of local abelian fields*, Acta Arith. **85** (1998), no. 3, 235–248. MR 1627831 (99d:11127)

[Ull70] S. Ullom, *Integral normal bases in Galois extensions of local fields*, Nagoya Math. J. **39** (1970), 141–148. MR 0263790 (41 #8390)

[Vin05] S. Vinatier, *Galois module structure in weakly ramified 3-extensions*, Acta Arith. **119** (2005), no. 2, 171–186. MR 2167720 (2006d:11135)

[Vos81] S. V. Vostokov, *Normal basis for an ideal in a local ring*, J. Sov. Math. **17** (1981), 1755–1758.

HENRI JOHNSTON, ST. JOHN'S COLLEGE, CAMBRIDGE CB2 1TP, UNITED KINGDOM
E-mail address: `H.Johnston@dpmms.cam.ac.uk`
URL: `http://www.dpmms.cam.ac.uk/~hlj31`